Buprenorphine Initiation Note
☐ Patient presents for in-clinic initiation
Evaluated using COW scale? □ Yes □ No
Scoredon COW Scale First Assessment
*If patient scored < 13 on COWS and/or reported illicit opioid use within 24 hours, do not proceed with in-clini initiation and consult OBAT provider for altered treatment plan including at home initiation.
Patient self-administeredmg sublingually as prescribed
<ul> <li>□ Assessed and instructed patient in proper administration</li> <li>□ Patient observed to tolerate medication</li> </ul>
Summary 1: COW Scale First Assessment
Resting Pulse Rate  □ 0 = pulse rate 80 bpm or below  □ 1 = pulse rate 80-100 bpm  □ 2 = pulse rate 101-120 bpm  □ 3 = pulse rate greater than 120 bpm
Sweating  □ 0 = no report of chills or flushing  □ 1 = subjective report of chills or flushing  □ 2 = flushed or observable moistness on face  □ 3 = beads of sweat on brow or face  □ 4 = sweat streaming off face
Restlessness during Assessment  □ 0 = able to sit still  □ 1 = reports difficulty sitting still, but is able to do so  □ 3 = frequent shifting or extraneous movements of legs/arms  □ 5 = unable to sit still for more than a few seconds
Pupil Size  □ 0 = pupils pinned or normal size for room light □ 1 = pupils possible larger than normal for room light □ 2 = pupils moderately dilated □ 5 = pupils so dilated that only the rim of the iris is visible
Bone or Joint Aches  □ 0 = not present □ 1 = mild diffuse discomfort

$\Box$ 2 = patient reports severe diffuse aching of joints/muscle $\Box$ 4 = patient is rubbing joints or muscles and is unable to sit still because of discomfort
Runny Nose or Tearing  □ 0 = not present  □ 1 = nasal stuffiness or unusually moist eyes  □ 2 = nose running or tearing  □ 4 = nose constantly running or tears streaming down cheeks
GI Upset  □ 0 = no GI symptoms  □ 1 = stomach cramps  □ 2 = nausea or loose stool  □ 3 = vomiting or diarrhea  □ 5 = multiple episodes of diarrhea or vomiting
Tremor  □ 0 = no tremor  □ 1 = tremor can be felt but not observed  □ 2 = slight tremor observable  □ 4 = gross tremor or muscle twitching
Yawning  □ 0 = no yawning  □ 1 = yawning once or twice during assessment  □ 2 = yawning three or more times during assessment  □ 4 = yawning several times per minute
Anxiety or Irritability  □ 0 = none □ 1 = patient reports increasing irritability or anxiousness □ 2 = patient obviously irritable/anxious □ 4 = patient so irritable or anxious that participation in the assessment is difficult
Gooseflesh Skin $\Box 0 = \text{skin is smooth}$ $\Box 3 = \text{piloerection of skin can be felt or hairs standing up on arms}$ $\Box 5 = \text{prominent piloerection}$
Total Score
Score: $5 - 12 = Mild$ 13 - 24 = Moderate 25 - 36 = Moderately Severe More than $36 = Severe Withdrawal$

☐ Patient presents for second initiation/re-evaluation
Evaluated using COWS scale? ☐ Yes ☐ No
Scoredon COWS Scale Second Assessment
Patient self-administeredmg sublingually as prescribed
☐ Assessed and instructed patient in proper administration☐ Patient observed to tolerate medication
☐ Patient presents for subsequent induction/re-evaluation
Evaluated using COWS scale? ☐ Yes ☐ No
Scoredon COWS Scale Second Assessment
Patient self-administeredmg sublingually as prescribed
☐ Assessed and instructed patient in proper administration ☐ Patient observed to tolerate medication

This document has been reproduced with permission from Boston Medical Center® and is excerpted from: Wason, K.F.; Potter, A.L.; Alves, J.D.; Loukas, V.L.; Lastimoso, C.S.; Sodder, S.B.; Caputo, A.M.; and LaBelle, C.T. *Massachusetts Nurse Care Manager Model of Office Based Addiction Treatment: Clinical Guidelines*. Unpublished treatment manual, Boston Medical Center, May 2021. This manual or any documents therein are not a substitute for informed medical decision making by an appropriate, licensed provider.